

DRIGALSKI GLUCOSE AGAR

A glucose containing selective and differential medium for detection and enumeration of Enterobacteriaceae.

Dehydrated media	
Code number:	500 g: DAC20500, 5 kg: DAC25000
Colour:	Beige
Appearance:	Homogeneous hygroscopic powder
pH before sterilization (25 °C):	7,1 – 7,5

Direction: Suspend **47 g** in one litre of distilled water and heat with frequent agitation until the medium boils well. Mix well before pouring.

Warning!

The medium is heat sensitive.
No further sterilisation is necessary or desirable.

Prepared media	
Bottled media:	100 ml: DAC30100, 500 ml: DAC30500
Plated media:	55 mm: DAC50055, 90 mm: DAC50090
Colour:	Green
pH (25 °C):	7,2 – 7,4

Direction: Dispense the melted bottled media aseptically into sterile Petri-dishes. Media in Petri-dishes are ready to use.

FORMULA in g/l

Peptones	18,400
Bile salt	1,500
Glucose	10,000
Sodium chloride	2,000
Bromothymol blue	0,150
Crystal violet	0,002
Agar	15,000

Note: The typical formula can be adjusted to obtain optimal performance.

Storage conditions: Store the dehydrated media tightly closed in a dry place at room temperature. Store the bottled media protected from light at room temperature. Store the plated media protected from light at 2-8 °C. Use before the expiry date on the label.

Quality control:

Test strains	Incubation temp: 37 °C	Growth	Incubation time: 24 h
<i>Escherichia coli</i>	ATCC 25922	Good, yellow colonies	
<i>Proteus mirabilis</i>	ATCC 29906	Good, yellow colonies without swarming	
<i>Enterococcus faecalis</i>	ATCC 29212	Inhibited	

References: Ewing (1986) Edwards and Ewing's identifications of the enterobacteriaceae, 4th ed.

In vitro diagnostic – for professional use only!