TECHNICAL SHEET



LETHEEN AGAR BASE

A highly nutritious medium that neutralizes quaternary ammonium compounds, for sampling of environmental surfaces that have been treated with disinfectants.

Dehydrated media			
Code number:	500 g: LTA20500, 5 kg: LTA25000		
Colour:	Yellowish		
Appearance:	Homogeneous hygroscopic powder		
pH before autoclaving (25 °C):	7,0 – 7,4		

Direction: Suspend **58 g** in one litre of distilled water. Add **5 ml of TWEEN 80 Supplement (TWS80100)**. Mix well and keep the suspension at about 50 °C until the lecithin dissolves completely (20 – 30 min). The dissolution is completed, when the medium is yellowish and slightly turbid, but exempt from any precipitate. Heat with frequent agitation until the medium boils well. Sterilise by autoclaving at 121 °C for 15 minutes.

Prepared media		
Bottled media:	100 ml: LTA30100, 500 ml: LTA30500	
Plated media:	55 mm: LTA50055, 90 mm: LTA50090 Contact Petri-dish: LTA50065	
Colour:	Yellowish, homogeneous turbid	
pH (25 °C):	7,1 – 7,3	

Direction: Dispense the melted bottled media aseptically into sterile Petri-dishes. Media in Petri-dishes are ready to use.

FORMULA in g/l

Peptones	32,2
Sodium chloride	5,0
Sodium bisulphite	0,1
Lecithin	0,7
Agar	20,0

Note: The typical formula can be adjusted to obtain optimal performance.

Storage conditions: Store the dehydrated media tightly closed in a dry place at room temperature. Store the bottled media protected from light at room temperature. Store the plated media protected from light at 2-8 °C. Use before the expiry date on the label.

Quality control:

Test strains	Incubation temp: 37 °C	Growth	Incubation time: 24 h
Staphylococcus aureus ATCC 29213		Good	

References: FDA (1992) Microbiological Methods for Cosmetics. Chapter 23.

In vitro diagnostic - for professional use only!

Valid from: 23 July 2016

Email: export@biolab.hu