

WL DIFFERENTIAL BROTH

A selective medium for the control of industrial fermentation, particularly the processing of beer.

Dehydrated media	
Code number:	500 g: WDB20500, 5 kg: WDB25000
Colour:	Beige
Appearance:	Homogeneous hygroscopic powder
pH before autoclaving (25 °C):	5,3 – 5,7

Direction: Suspend **60 g** in one litre of distilled water and heat gently to dissolve the medium completely. Dispense into final containers and sterilise by autoclaving at 121 °C for 15 minutes.

Prepared media	
Bottled media:	100 ml: WDB30100, 500 ml: WDB30500
Tubed media:	150 x 15 mm: WDB40010 (10 ml)
Colour:	Greenish
pH (25 °C):	5,4 – 5,6

Direction: Dispense the bottled media aseptically into sterile final containers. Media in tubes are ready to use.

FORMULA in g/l

Peptones	9,2000
Glucose	50,0000
Potassium chloride	0,5500
Calcium chloride	0,1250
Magnesium sulphate	0,1250
Ferric chloride	0,0025
Manganese sulphate	0,0025
Cycloheximide	0,0040
Bromocresol green	0,0220

Note: The typical formula can be adjusted to obtain optimal performance.

Storage conditions: Store the dehydrated media tightly closed in a dry place at room temperature. Store the bottled and tubed media protected from light at room temperature. Use before the expiry date on the label.

Quality control:

Test strains	Incubation temp: 30 °C	Growth	Incubation time: 48 h
<i>Lactobacillus fermentum</i> ATCC 9338		Good	
<i>Saccharomyces cerevisiae</i> ATCC 9763		Inhibited	

References: Green and Gray (1950) Wallerstein Lab. Commun. 13: 357.

In vitro diagnostic – for professional use only!